

HAIR RESTORATION


CLINICAL STUDY TO EVALUATE THE EFFICACY OF A COMBINED TREATMENT WITH XL HAIR® IN DIFFERENT BALDNESS PATTERNS

E Ranneva (Spain); G Siquier (Netherlands); Ph Deprez (Spain); R Vergereau (France).
2011-12-13

INTRODUCTION

Hair loss becomes one of the most frequent problems of both genders, which motivates the patients to visit the dermatologist. Male and female baldness patterns have statistical, clinical and diagnostic differences. The medical treatments FDA approved for hair loss include corticosteroids, minoxidil and 5-alpha-reductase inhibitors. The surgical treatments for hair restoration are strip techniques, follicular unit extraction, scalp flap and scalp reduction. Each product or method have contraindications and side effects.


XL Hair® injectable class III medical device, **Reparestim® Hair TD** and **AD Daily Care Hair** are capillary regenerators which revitalize and strengthen capillary fibre via a greater contribution of essential nutrients for capillary growth and a stimulating action of hair growth factors.



MATERIAL AND METHODS

A multi-center, open label, non-comparative pilot study was performed in 47 patients (32 women and 15 men) who had medical visit due to hair loss. The average age of the patients was 34 years and the evolution period of the alopecia condition was 5 years. Efficacy was determined at the 3rd and 6th months.

XL Hair® injections and/or **Reparestim® Hair TD** application with microneedling device **AD Roll TD®** repeated once a week on the area of hair loss or alopecia, total 8 sessions. Possible side effects were easily treated, as daily care **AD Daily Care Hair** spray has been applied directly on the scalp with light massage ensuring the product is well distributed over the problematic area. Application twice a day, 3 times a week, during 8-16 weeks. The regular treatment with injections of **XL Hair®** or microneedling with **Reparestim® Hair TD** combined with topical application of **AD Daily Care Hair** could stop active hair loss in 3 months and stimulate new hair growth in 6 months. The best efficacy was observed in case of symptomatic hair loss in the female pattern: after pregnancy, nursing period, stress, starvation. In the male pattern which is related to the androgenic alopecia, the result gives satisfaction, but further investigation will be necessary. The patient with androgenic alopecia received 12 treatments, once every two weeks. 60 % of the patients were satisfied with result of the treatment after 12 weeks.



CONCLUSION

The results of the study indicate that injections of **XL Hair®** or **Reparestim® Hair TD** applied with microneedling device (**AD Roll TD®**) and combined with **AD Daily Care Hair** are efficient treatment for hair loss in different baldness patterns. The results indicate that mesotherapy injections of **XL Hair®**, or microneedling with **Reparestim® Hair TD** combined with **AD Daily Care Hair**: induce the activation of the hair follicle which promotes enlargement of the anagen phase and a shortening of the telogen phase, reversing the miniaturization of the hair follicles besides stop hair loss and promote new hair growth. The protocol of application and frequency of treatment have been adapted to the gender. To optimize obtained results, the application of **AD Daily Care Hair** cannot be less than 6 months.

Thanks to Aesthetic Dermal S.L. for its support on the clinical research.

aestheticdermal.com - **rrs-inject.com**

For any supplementary information: info@aestheticdermal.com